

# Heterosis of Early Growth Performance in Three Breeds of Rabbits (*Oryctolagus cuniculus*)

**Nwakpu, P. E.**

Department of Animal Science,  
Ebonyi State University, Abakaliki  
Email: penwakpu.380@gmail.com

**Ogbu, C. C.**

Department of Animal Science,  
MOUA, Umudike, Umuahia

**Ugwu S. O. C.**

Department of Animal Science,  
University of Nigeria, Nsukka

**Abstract** – The study investigated heterosis effects for early growth performance using three popular breeds of rabbits in Nigeria (Chinchilla, New Zealand White and Dutchbreeds). Thirty six (36) sexually mature rabbits belonging to three genetic groups (Twelve per breed at the mating ratio of three males to nine females) were employed for the study. Matings within each breed were carried out to generate purebred progenies. These were reared and monitored for growth performance from birth to 32 weeks of age (age of sexual maturity). At 32 weeks of age, 3 males and 9 females each of the three breeds were randomly selected from their respective groups and used in cross breeding in a ratio of one Buck to two does. F<sub>1</sub> crossbred progenies were reared according to crossbred groups to weaning (8 weeks of age). The bioeconomic traits considered in this investigation for the purebreds and F<sub>1</sub> crossbred groups were; individual kitten weight at birth and at bi-weekly intervals up to 8 weeks of age; individual kitten weight at weaning; litter size at birth and at weaning sex-ratio, pre-weaning mortality and estimates of heterosis. Data collected were subjected to Analysis of Variance to test for the effect of breeding groups on parameters measured. Comparison between crossbred and purebred groups was done using independent samples (T-Test), while significance of heterosis was performed using one-sample T-Test. All analyses was performed using the Statistical Package for Social Sciences (SPSS) Version 17.0 (2007). Significant means were separated using the Duncans New Multiple Range Test in SPSS. Results showed that the crossbred kits were heavier in weight (g) than the purebred kits from birth to weaning (45.39g from Nzw x Nzw to 48.35g from CH x CH among the purebreds and 38.25g from Nzw x CH to 55.50g from D x CH) for the crossbreds at birth; and rose to 420.14g from D x D and 569.38g from CH x CH among the purebreds to 670.63g from Nzw x CH and 704.41g from Nzw x D) for the crossbreds. The result of the reproductive performance of the breeds indicated lack of significant (P>0.05) differences except in the preweaning mortality of the purebreds and the litter size at weaning (LSW) among the crossbreds. Litter size at birth (LSB) were (on average) lower for the purebreds than the crossbreds. Estimates of direct heterosis for body weight were all positive and favorable for the crossbred kits except (Nzw x CH) crossbred kits. Positive heterosis for litter size at birth ranged from 14% to 26% and 14% to 50% for litter size at weaning among the crossbred kits. Within litter mortality in Nzw x D and Nzw x CH crosses were negative but increased and were positive among the D x CH crossbred kits. The potential of crossbreeding to tap the non-additive genes for reproductive performance was confirmed in the present study. Crossing (Nzw x CH) and (Nzw x D) resulted in positive heterosis for body weight, while (D x CH) helped to improve preweaning mortality and litter size.

**Keywords** – Growth Performance, Breeds of Rabbits, Heterosis, Genetic Groups, T-Test.

## I. INTRODUCTION

Growth traits in growing rabbits are important because heavier marketable body weight promotes the economics of rabbit production (1). Cross breeding is one of the fast tools offered to breeders to improve many traits in farm animals (2). One major basis for crossbreeding lies in the fact that the crossbred animals often manifest heterosis due to good interbreeding or nicking ability.

Heterosis usually occurs where the populations crossed differ genetically. Positive heterosis is the superiority of the heterozygous above the average value of the two parents. The percent superiority depends on the magnitude of non-additive genetic variation in the trait under consideration (3). Breed complementation for growth and carcass traits by crossing animals of different genotypes has generated a lot of interest among rabbit breeders (4), (5), (6), (7), (8), (9). For instance, breed complementary was achieved by crossing the fast growing indigenous Sinar Gabali breed with the imported V-rabbit line (9). Also similar result by (10) was observed in a study to investigate the heterosis effects for preweaning litter characteristics using three Nigerian local rabbits (local white, local brown tail and local black). The needed information on breed complementarity which should precede commercial trend is much less documented in developing countries like Nigeria.

Commercial rabbit production in Nigeria has been maintained mainly through the use of imported exotic breeds and their crosses. Consequently, the availability of breeds differing in their growth characteristics is an important source of flexibility that can allow more rapid adjustments to changing economic conditions in the country. However, a sound knowledge of this important aspect of animal production and its interaction with the environment is essential for a maximum and rational utilization of the breeding stock.

Reference (11) had observed that, when crosses are made between two parental lines which differ genetically, such as different inbred lines or different breeds or strains within a particular specie, heterozygosity is presumably generated in the resulting F<sub>1</sub> progeny. The mean performance of the F<sub>1</sub> progeny may exceed the mean of both parents (mid-parents heterosis) or the performance of the poorer parents (Poor parent heterosis). In the same vein, (12) also stated that the genetic explanation for the hybrid extra vigor is basically the same, whether it be animal or crop. Heterosis is produced by the fact that, the dominant gene of a parent is usually more favorable than the recessive partner. When the genetic group differ in

frequency of gene they have and dominance exist, then heterosis will be produced.

In animal breeding, crossbreeding allows the exploitation of breed qualities from genetic and biological point of view (13). It also enables breed discrimination and the analysis of genetic variability. To effectively utilize genetic resources, the genetic and environmental causes of phenotypic variation in economic attributes need to be differentiated (14). Furthermore, the genetic components have to be decomposed into direct and maternal effects (15). Characterization of genetic and maternal effects attributable to each breed or breed combinations enable producers to choose amongst breeds and to assign appropriate role(s) to selected breeds in breeding schemes.

Crossing different breeds may lead to improved performance over the pure/inbred parents. This improved performance is recognized as heterosis. In secondary generations such as backcrosses, epistatic effects may also be important. Although considerable research has evaluated cross breeding in farm animals in Nigeria, the analysis and reporting of the results have often been inadequate. Most reports presented only the breed group means and mid-parent heterosis values paying little or no attention to the evaluation of the genetic components: direct additive and maternal effects, direct heterosis, maternal heterosis, recombination loss, over dominance, relative heterosis, heterobeltiosis etc, or to the development of parameters that allow prediction of the performance of crosses that have not actually been tested (16).

Accordingly, this study is designed to estimate the crossbreeding parameters of the main economically important traits in cross involving three breeds commonly reared in Nigeria. This work shall investigate the breed additive effects (direct and maternal) and heterosis effects on growth, litter size traits, and preweaning mortality rate of the breeds.

### *1.1 Objectives of the Study*

The main objective of this investigation is to determine the heterosis of early growth performance and other reproductive traits of economic importance in three popular breeds of rabbits. Other objectives include;

- a) To determine the heterosis involved in traits like litter size at weaning (LSW), litter size at birth (LSB), and preweaning mortalities among the three popular breeds of rabbits in Nigeria.
- b) This study shall also ascertain heterosis for litter weight at birth and weaning in order to establish the impact of crossbreeding on these traits.

### *1.2 Justification of the Study*

One major advantage of cross breeding is breed complementarity. This exercise will help to expose the potential of crossbreeding in helping to tap the non-additive genes for possible improvement in the reproductive performance. For instance, (17) observed a positive heterosis of 28% for litter weight in a synthetic breed (APRI) when they crossed highly adaptable Egyptian Baladi Red with the Spanish V-line noted for prolificacy and mothering ability.

A study by (18) showed a less successful result (92.5% to 5.0% for total gain and 0.7% to 9.5% for daily gain) in the local Baladi Red x New Zealand white crossed kits.

The variations in the manifestations of heterosis indicate that in some cases, heterosis is apparent at birth, while in others, it does not appear until a considerable period following birth. Among rabbit crosses made by previous workers, only one cross, polish and Himalayan (19) produced F<sub>1</sub> rabbits which were heavier than both parental races. The body weight of the F<sub>1</sub> rabbits in castle's experiment was 45% heavier than the polish and 9% heavier than the Himalayan. However, (17) crossed Flemish and Ermine, and White Giant and Ermine rabbits without heterosis in the F<sub>1</sub> generation, but observed heterosis in haemoglobin value, erythrocyte number and viability. It was probable that the extremely heavy rabbits e.g. Flemish, had a larger number of effective quantitative growth genes than the extremely small rabbits e.g. Polish and that F<sub>1</sub> from extremely heavy and extremely small rabbits received half the effective growth genes from each parent, and became intermediate in size. Nevertheless, the size of the F<sub>1</sub> rabbits was closer to that of the heavy parent. This probably was due to the influence of heterosis derived from certain pairs of effective allelic growth genes. Most of these genes were probably paired in the cross and produced heterosis which made the F<sub>1</sub> animals heavier than both parental strains.

In meat rabbit industry, there is need to develop sire breeds and (or) lines to cross to available dam breeds to potentially enhance breed complementarity and heterosis benefits on post weaning fryer performance. Commercial rabbit meat is usually produced by a three-way cross involving crossbred females mated to males from a sire line. The crossbred females are obtained by mating males and females from two female lines selected for litter size, while the sire lines are generally selected for growth rate, carcass yield and meat quality (20), (21). When designing a genetic selection programme, two aspects must be considered; the choice of partner lines and their crossbreeding scheme and the within-line selection. Concerning the choice of partner lines and their optimal utilization, different crossbreeding schemes have to be tested in order to identify or to predict the best combination of line. (22) has theorized the choice of breeds and breed combinations and proposed a set of parameters allowing the prediction of performance from a potential breed combination.

This study shall aim at comparing different crosses of three rabbit breeds (New Zealand White, Dutch and Chinchilla) and estimate heterosis and direct and maternal additive effects, as well as some non-genetic effects (litter size, mortality) on rabbit growth, so as to identify the best crossbreeding plan to use for rabbit meat production in the south eastern derived savanna zone of Nigeria.

## **II. MATERIALS AND METHODS**

### *2.1 Experimental Site*

The breeding bucks and does used for this experiment were raised at the Teaching and Research Unit of the

Department of Animal Science, Ebonyi State University, Abakaliki.

### 2.2 Experimental Animals

Thirty six (36) sexually mature rabbits belonging to three genetic groups (twelve per breed at the mating ratio of three males to nine females) were employed for the study. Mating within each breed was carried out to generate purebred progenies. These were reared and monitored for growth performance from birth to 32 weeks of age (age at sexual maturity). At 32 weeks of age, three males and nine females each of New Zealand white, Dutch, and Chinchilla were randomly selected from their respective groups and used in crossbreeding according to the following scheme: New Zealand white (NZW) female with Dutch (D) male (NZW x D); New Zealand white female with Chinchilla (CH) male (NZW x CH); and Dutch female with Chinchilla male (D x CH) in a ratio of one buck: two does. F<sub>1</sub> crossbred progenies were reared according to crossbred groups to weaning at 8 weeks of age.

### 2.3 Management of the Animals

The three breeds of rabbits used were sourced from a reputable farm (Emmpina Farms, Akwa Ibom State). They were housed in hutches measuring about 75cm x 63m x 54cm. Kits and Does were fed mash diet containing 46.59% rice bran, 25.00% palm kernel cake, 15.50% wheat offal, 1.50% Oyster shell, 0.55% bone meal and 0.5% salt. The mash diet was supplemented with available local forages (Banana leaves, Centrosema Pubescence). The experiment was carried out under a hygienic environment by sweeping the rabbitry unit and cages thoroughly every day and disinfected weekly using concentrated isol solution. The animals were injected with ivermectin for the treatment of both ecto and endoparasites at bi-monthly intervals.

### 2.4 Data Collection

The bio-economic traits considered in this investigation for the purebreds and F<sub>1</sub> crossbred groups are:

- Individual kitten weight at birth and at bi-weekly intervals up to 8 weeks of age.
- Individual kitten weight at weaning (8 weeks of age)
- Litter size at birth
- Litter size at weaning
- Number of males and females at birth and at weaning.
- Pre-weaning mortality (%)
- Sex-ratio at birth and at weaning
- Estimates of direct heteroses were obtained following (Dickerson, 1992)

$$H^1 (NZW \times D) = \{NZW \times D\} - \{NZW \times NZW\} \{D \times D\}$$

$$H^1 (NZW \times CH) = \{NZW \times CH\} - \{NZW \times CH\} \{NZW \times NZW\} \{CH \times CH\}$$

$$H^1 (D \times CH) = \{D \times CH\} - \{D \times D\} \{CH \times CH\}$$

NZW, D and CH represent the New Zealand white, Dutch and Chinchilla breeds respectively. The percent heterosis shall be collected as follows:

$$\frac{H^1 (\text{Units})}{\text{Means of purebreds}} \times \frac{100}{1}$$

H<sup>1</sup> = Estimate of heterosis in units

### 2.5 Data Analysis

Data collected were subjected to analysis of variance to test for effect of breeding groups on parameters measured. Comparison between crossbred and purebred groups was done using independent samples T-Test, while significance of heterosis was performed using one-sample T-Test. All analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 17.0 (2007). Significant means were separated using the Duncan New Multiple Range Test in SPSS.

The statistical model used in this study is as follows:-

$$Y_{ij} = \mu + B_i + e_{ij}, \text{ where}$$

Y<sub>ij</sub> = Record of the jth individual or litter belonging to the ith breed of the rabbit,

μ = Random mean,

B<sub>i</sub> = Effect of the ith breed

e<sub>ij</sub> = Residual error

## III. RESULTS

Table 1: Mean ( $\pm$  S.E) of the body weight (g) performance of the three breeds of rabbits and their F<sub>1</sub> crosses.

Age (Wks)	Purebred			F <sub>1</sub> Crossbred		
	CH x CH	NZW x NZW	D x D	D x CH	NZW x CH	NZW x D
0	48.35 $\pm$ 1.05	45.39 $\pm$ 1.06	45.40 $\pm$ 1.12	55.5 $\pm$ 0.73 <sup>a</sup>	38.25 $\pm$ 0.71 <sup>b</sup>	53.84 $\pm$ 0.66 <sup>b</sup>
2	156.00 $\pm$ 3.19 <sup>a</sup>	155.39 $\pm$ 3.23 <sup>a</sup>	137.14 $\pm$ 3.41 <sup>b</sup>	156.43 $\pm$ 1.31	159.59 $\pm$ 1.27	160.05 $\pm$ 1.18
4	287.13 $\pm$ 5.89 <sup>a</sup>	241.03 $\pm$ 5.96 <sup>b</sup>	212.43 $\pm$ 6.30 <sup>c</sup>	374.47 $\pm$ 3.31 <sup>b</sup>	342.41 $\pm$ 3.21 <sup>c</sup>	408.92 $\pm$ 2.98 <sup>a</sup>
6	398.75 $\pm$ 10.23 <sup>a</sup>	281.15 $\pm$ 10.36 <sup>b</sup>	278.29 $\pm$ 10.94 <sup>b</sup>	551.07 $\pm$ 6.54 <sup>a</sup>	501.81 $\pm$ 6.33 <sup>b</sup>	506.84 $\pm$ 5.89 <sup>b</sup>
8	569.38 $\pm$ 10.90 <sup>a</sup>	435.13 $\pm$ 11.04 <sup>b</sup>	420.14 $\pm$ 11.65 <sup>b</sup>	696.10 $\pm$ 6.42 <sup>a</sup>	670.63 $\pm$ 6.21 <sup>b</sup>	704.41 $\pm$ 5.78 <sup>a</sup>

a, b, c: Means not followed by the same superscripts are significantly (P>0.05) different for the purebred and crossbreds.

Data on the body weight (g) performance of the three breeds of purebred and their F<sub>1</sub> crosses of rabbits were presented in Table 1. Result indicated lack of significant (P>0.05) differences in the body weight of the purebreds

at birth but varied significantly (P<0.05) as from 2<sup>nd</sup> week of birth up to their 8<sup>th</sup> week.

However, their F<sub>1</sub> crossbreds differed significantly (P<0.05) as from birth but did not vary at their 2<sup>nd</sup> week of age. Significant variation (P<0.05) in body weight was

recorded as from the 4<sup>th</sup> week of age continuously up to their point of weaning at 8<sup>th</sup> week of age.

The Chinchilla breed were the heaviest in body weight among the purebreds, followed by the New Zealand white and the least in weights were the Dutch breed. In the F<sub>1</sub> crossbreds, the cross between the male Dutch and Female Chinchilla were the heaviest in weight followed by the crossbreds from male New Zealand and female Dutch

although not statistically ( $P > 0.05$ ) different from the cross between New Zealand male and female Chinchilla at birth. This trend however dropped in week 2, 4 and 8 in favour of the cross between the male New Zealand white and Dutch female as the animals approached their weaning period at 8 in favour of the cross between the male New Zealand white and Dutch female as the animals approached their weaning period at 8 week of age.

Table 2: Mean ( $\pm$ S.E) of the reproductive traits of the purebred and their F<sub>1</sub> crosses

Traits	Purebred			F <sub>1</sub> Crossbred		
	CH x CH	NZW x NZW	D x D	D x CH	NZW x CH	NZW x D
LSB	5.33 $\pm$ 0.44	5.78 $\pm$ 0.40	5.38 $\pm$ 0.38	6.17 $\pm$ 0.40	6.50 $\pm$ 0.34	7.17 $\pm$ 0.31
LSW	4.44 $\pm$ 0.24	4.33 $\pm$ 0.24	4.25 $\pm$ 0.37	4.83 $\pm$ 0.40 <sup>b</sup>	5.33 $\pm$ 0.21 <sup>ab</sup>	6.17 $\pm$ 0.48 <sup>a</sup>
PWM	14.68 $\pm$ 3.10 <sup>b</sup>	20.56 $\pm$ 4.55 <sup>a</sup>	19.17 $\pm$ 6.63 <sup>a</sup>	20.95 $\pm$ 5.84	16.97 $\pm$ 4.89	13.69 $\pm$ 6.4
MAB	2.13 $\pm$ 0.10	2.51 $\pm$ 0.14	2.85 $\pm$ 0.14	3.50 $\pm$ 0.49	3.17 $\pm$ 0.49	4.33 $\pm$ 0.49
FAB	2.05 $\pm$ 0.11	2.62 $\pm$ 0.12	2.10 $\pm$ 0.08	2.67 $\pm$ 0.44	3.33 $\pm$ 0.44	2.83 $\pm$ 0.44
MAW	1.80 $\pm$ 0.13	2.0 $\pm$ 0.20	2.3 $\pm$ 0.13	2.50 $\pm$ 0.40	2.67 $\pm$ 0.40	3.33 $\pm$ 0.40
FAW	1.90 $\pm$ 0.17	2.10 $\pm$ 0.18	2.30 $\pm$ 0.05	2.33 $\pm$ 0.36	2.67 $\pm$ 0.36	2.83 $\pm$ 0.36
SRB	0.43 $\pm$ 0.03	0.33 $\pm$ 0.01	0.42 $\pm$ 0.05	0.56 $\pm$ 0.06	0.49 $\pm$ 0.06	0.60 $\pm$ 0.06
SRW	0.31 $\pm$ 0.07	0.41 $\pm$ 0.07	0.45 $\pm$ 0.05	0.52 $\pm$ 0.06	0.51 $\pm$ 0.06	0.53 $\pm$ 0.06

a, b: Means not followed by same superscript on the same row are statistically ( $P < 0.05$ ) different.

LSB: Litter size at birth, LSW: Litter size at weaning, PWM: Pre-weaning mortality, MAB: Males at birth, FAB: Females at birth, MAW: Males at weaning, FAW: Females at weaning, SRB: Sex-ratio at birth SRW: Sex-ratio at weaning

Table 2 presents the results of data on litter size at birth, litter size at weaning, post weaning mortality rate, males and females at birth and at weaning as well as the sex ratio at birth and at weaning. The pre-weaning mortality (PWM) varied from 14.68% (CH x CH) to 20.56% (NZW x NZW) which also did not differ from 19.17% (D x D)

purebreds. There was lack of significant ( $P > 0.05$ ) differences in the reproductive traits measured among the three popular breeds of rabbits used in the experiment. Similar trend as in the purebreds was observed among the crossbreds except in the litter size at weaning (LSW) that showed significant ( $P < 0.05$ ) differences and ranged from 4.83 (D x CH) to 6.17 (NZW x D) per litter. Litter sizes at birth and at weaning were higher numerically among the crossbreds than the purebreds. Pre-weaning survivability in the crossbreds was highest in the New Zealand white males crossbred with Dutch female crossbreds.

Table 3: Estimates of direct heterosis (%) for body weight (g) performance of three breeds of rabbits

CROSSBREDS			
Age (weeks)	NZW x D	NZW x CH	D x CH
0	25.06 $\pm$ 2.59 **	-14.09 $\pm$ 2.55 **	22.46 $\pm$ 3.57 **
2	10.81 $\pm$ 3.44 **	05.09 $\pm$ 3.39 **	10.30 $\pm$ 4.73 **
4	79.19 $\pm$ 4.07 **	25.65 $\pm$ 4.01 **	49.14 $\pm$ 5.60 **
6	75.81 $\pm$ 4.37 **	44.36 $\pm$ 4.31 **	56.73 $\pm$ 6.02 **
8	61.38 $\pm$ 4.55 **	33.16 $\pm$ 4.48 **	41.89 $\pm$ 6.26 **

\*\*Heteroseres were highly significant ( $P < 0.01$ ) among the crossbreds.

The percent heterosis effects for body weight were positive in NZW x D and D x CH crossbred kits and negative for NZW x CH crossbred kits ranging from (+25.06%, +22.46% and -14.09%) respectively at birth. Percent heteroseres were however positive for body weight (g) from birth to 8<sup>th</sup> week (point of weaning) for all the

crossbred kits. The highest positive heterosis of 79.19% was recorded from the NZW x D crossbred kits at 4<sup>th</sup> week and decreased gradually from the 6<sup>th</sup> and 8<sup>th</sup> week of age respectively. Percent heterosis effects were highly significantly ( $P < 0.01$ ) different among the crossbreds from birth to weaning (Table 3).

**Table 4: Heterosis (%) of litter traits for various breeding groups**

Groups	LITTER TRAITS		
	LSB	LSW	PW
Breeding groups			
NZW x D	0.26±0.17 <sup>NS</sup>	0.50±0.17 <sup>NS</sup>	-0.25±0.18 <sup>NS</sup>
NZW x CH	0.17±0.14 <sup>NS</sup>	0.26±0.14 <sup>NS</sup>	-0.18±0.15 <sup>NS</sup>
D x CH	0.14±0.30 <sup>NS</sup>	0.14±0.30 <sup>NS</sup>	0.17±0.33 <sup>NS</sup>

NS: Non-significant differences (P>0.05) in the litter trait heteroses.

Table 4 presents the percent heterosis effects for litter traits among the crossbred kits. Result showed non-significant (P>0.05) differences in the heteroses of litter traits of the crossbred kits. Percent heterosis were negative for the pre-weaning mortality in NZW x D and NZW x CH crossbreds and positive for D x CH crossbreds. Highest percent heterosis of litter size at weaning was recorded in the NZW x D crossbred kits and lowest for the D x CH crossbred kits.

#### IV. DISCUSSION

##### 4.1 Growth Performance of the Breeds

Data on the body weight (g) of the three breeds of the rabbits (purebreds and their F<sub>1</sub> crosses) are presented in Table 1. Mean body weight (g) varied from 45.39 (NZW x NZW) to 48.35 (CH x CH) among the purebreds and 38.25 (NZW x CH) to 55.5 (D x D) for the crossbreds at birth and rose to 420.14 (D x D) and 569.38 (CH x CH) among the purebreds to 670.63 (NZW x CH) and 704.41 (NZW x D) for the crossbreds. Crossbred kits in this study were heavier than the purebreds kits from birth to weaning at 8 week of age due to the smaller litters in which they were raised. Previous studies by (10), (23) and (1) showed that kit weight at birth and weaning body weight as well as post weaning body weight and market age were significantly higher for crossbred breeds or strains of rabbits. The present study disagrees with the report of (24) who observed a decline in growth rate with advancing kits age in their work on New Zealand white, Californian and Gray Giant Flander breeds. Other rabbit crosses made by previous workers showed that only one cross, polish and Himalayan (19) produced F<sub>1</sub> rabbits which were heavier than both parental races. The body weight of the F<sub>1</sub> rabbits in castle's experiment was 45% heavier than the polish and 9% heavier than the Himalayan. The differences in body weight of the crossbreds over the purebreds in this work did not reach the levels recorded in castle's work.

##### 4.2 Reproductive Performance

The results of the reproductive performance of the purebreds and their F<sub>1</sub> crossbreds are presented in table 2. Results indicated lack of significant (P>0.05) differences in the reproductive traits measured except that the preweaning mortality in the present study was higher compared with 5-7% of still born reported for highly prolific does in modern French Rabbitries (10) and (6). The present study also agrees with the submission of (8) who observed that about 94% of preweaning losses occurred within the first two weeks of life.

Litter size at birth were (on average) lower for the purebreds than the crossbreds. Generally, all the

reproductive traits measured except the preweaning mortality were numerically (on the average) higher among the crossbreds than the purebreds. The pre weaning survivability of the rabbits was also higher among the crossbreds than the purebred.

Litter size at birth and at weaning as well as the sex ratio at birth and at weaning recorded in this study was similar to that reported by (25), for Sudanese local rabbit but lower compared with litter size at birth in Egyptian and Exotic rabbits (26, 1).

First parity litter size at birth (total number born or total number born alive has been the selection criterion most frequently adopted to modify litter size genetically with no attention given to litter size at subsequent parities which has a major environment component influencing the litter size (8).

The differences in litter traits at birth due to parity effect may be attributed to changes in the physiological efficiency such as those associated with ovulation rate, ova wastage, implantation, embryo survival rate, embryonic mortality, uterine capacity, efficiency of providing the young with nourishment and intra uterine environment during the pre-natal development and also may be due to difference on the age of doe within parity (18).

However, the basis of the production is the genotype but the manifestation of genetic depends on numerous elements from the moment of fertilization; the surrounding in the uterus (space and blood supply of fetuses), the milk supply of kits, the nutritional and environmental condition during rearing, age at first mating, the physiological status at mating etc; can modify the performance of does; this implies that the reproductive performance of rabbit does is influenced by several genetic; physiological and environmental factors (27).

##### 4.3 Heterosis for Preweaning Body Weight

Estimates of direct heterosis for body weights were all positive and favorable for the crossbred kits in this study (Table 3) except (NZW x CH) crossbred kits at birth which was negative (-14.09%). Heterosis effect for body weight at birth was highest (25.06%) for (NZW x D) crossbreds and (22.46%) for (D x CH) crossbreds respectively at birth, dropped by more than fifty percent at the second week of birth and attained the highest value at week four (79.19%) before dropping gradually as they animals approached their weaning age of 8 week.

There were fluctuations in the bi-weekly heterosis effect for body weight among the rabbit crosses. Crossbreeding was unfavourable for growth rate in NZW x CH crossbred kits at birth and the 2<sup>nd</sup> week postpartum. Generally, crossbreeding assisted tremendously in improving the

growth rate of crossbred rabbits up to weaning and even beyond.

Previous works by (10), (28), (29), (30) revealed significant heterosis for growth rate and bodyweight, though at variable values of the parental means. Reference (31) observed that estimates of direct heterosis were and ranged from 4.9 to 16.7% for body weight; but the estimates for maternal heterosis were in most cases significantly negative (-4.5 to -15.2%) for body weight. Estimates of heterosis effect obtained by (24) for body weight and relative growth rate in post weaned New Zealand white, California and Flander crossbred rabbits were generally negative. On the other hand, a high positive heterosis effect (42.27%) for rate of growth at 10-12 weeks of age was observed in California x Flander rabbits.

In animal breeding, crossbreeding allows the exploitation of breed qualities from genetic and biological point of view (13). And one major advantage of crossbreeding is breed complementarity. This was demonstrated in a synthetic breed which combined the adaptability of Egyptian Baladi Red with prolificacy and mothering ability of the Spanish V-line. The synthetic breed showed a direct heterosis of 28% for litter weight (17). The results of this work have shown far higher heterosis in the bodyweight of the crossbred kits than the synthetic breed. A study by (18) showed a less successful result in the Local Baladi Red and New Zealand white crossbred kits.

#### 4.4 Heterosis for Prewaning Litter Size

There was a non-significant ( $P < 0.05$ ) differences in the heterosis for the preweaning litter size traits. The percent heterosis effects were positive among the crossbred kits from birth to weaning age. A positive direct heterosis for litter size at birth ranged from 14% to 26% and 14% to 50% for litter size at weaning among the crossbred kits.

The differences in heterosis effect for litter size among breed crosses in this study is in line with previous studies on New Zealand white and local Egyptian rabbits (1). Breed combination to improve litter size has generated a lot of interest among rabbit breeders (8), (9), (10). For instance, a new line named APRI, Produced by (17) showed a direct heterosis of 24 and 28 percent for litter size at birth and at weaning.

In a contemporary study, (28) observed a positive heterosis for litter size at birth and at weaning in their four-breed crosses under Egyptian rearing conditions. Reference (32) evaluated traits such as litter size at birth and at weaning in Spanish V-line x Saudi Gabali rabbits and reported that direct heterosis varied from 5.3% to 27.5% among the investigated production traits. However, the crossing of Sinai Gabali breed with the Spanish V-line did not give a significant individual heterosis for litter size at birth and at weaning (9).

#### 4.5 Heterosis for Prewaning Mortality

Result of this experiment demonstrate strongly that crossbreeding can help to reduce pre-weaning mortality among crossbred kits. Within litter mortality in NZW x D and NZW x CH crosses were negative but increased and were positive in D x CH crossbred kits (Table 4). The present work agrees with the earlier findings of (1),

(32) Both workers reported that preweaning mortality (PWM) was reduced when New Zealand rabbit was crossed with local Egyptian rabbit in a crossbreeding programme.

The superiority of the crossbred over the purebred in this work suggests that there exist a quantum of non-additive genetic effects to improve kits liveability especially in NZW x D and NZW x CH crossbred rabbits (3).

## V. CONCLUSION

The potential of crossbreeding to tap the non-additive genes for improving the reproductive performance are confirmed in the present study. (Nzw x CH) and (Nzw x D) crossbred kits had a positive heterosis for body weight, while (D x CH) was superior in pre-weaning mortality and litter size.

## REFERENCES

- [1] A.A. Rashwan, K.A Yaman; and A.I Abd El-Ghani (1997). "Performance of 3 Rabbit Strains and their Reciprocal Crosses in Egypt during Summer". *World Rabbit Science* Vol. 2 pp 55-60
- [2] S. O. Odubote, I. O. Akinokun and B. Somade (1997). "Productivity Levels of Three Breeds of Rabbits and Their Cross over a Three Year period in the Humid Tropics". *Arch Tierz* Vol. 40 pp 469-76.
- [3] N.S. Nagpure, M.D. Kotbber, A.K. Gore and S.N. Des Hunkh (1991). "Estimation of Heterosis in Crosses of Four Breeds of Rabbits". *Journal of Applied Rabbit Research*, Vol. 14, pp. 34-37.
- [4] R.D. Carregal, and J.F Lin (1984). "Effect of Mother Ability and General Combining Ability in Cross Rabbit (in Portuguese)" *Rev Soc. Bras Zootech.* Vol. 13 No 4 pp. 440.
- [5] M. Panic, M. Petkovic, and M. Urosenic, (1985). "The Most Useful Genetic Combinations of Rabbits for Broiler Production (in Polish)" *Vet. Glas.* Vol. 39 No. 3 pp 285.
- [6] F. Lebas, P. Coudert, R. Rouvier, and H. de Rochambeau (1986). *The Rabbit; Husbandry, Health and Production*, FAO, Rome, Italy.
- [7] C.E. Ozimba and S.D. Lukefahr (1991). "Evaluation of Purebred and Crossbred Rabbits for Carcass Merits". *Journal of Animal Science* Vol. 69, pp 2371.
- [8] A.A Rashwan and I.F.M. Marai (2000). "Mortality in Young Rabbits; A Review". *World Rabbit Science* Vol. 8 No. 3 pp 111-124.
- [9] M.M. Iraqi, E.A. Afifi, M. Baselga, M.H. Khalil, and M.L. Garcia (2008). "Additive and Heterotic Components for Post-weaning Growth Traits in a Crossing Project of V-line with Gabali rabbits in Egypt". *Proceedings of the 9<sup>th</sup> World Rabbit Congress, Verona, Italy*, pp 131-135.
- [10] T.R. Fayeye, (2013). "Heterosis for Prewaning Litter Characteristics in Nigerian Rabbit Crosses. *International Journal of Agricultural and Food Research*, vol. 2 No. 2 pp 1-6
- [11] S.N. Ibe, (1998). *Improved Genetic Stock; Nigerian Poultry Industry in the 1990s.* in Carddosa vol. 1 No. 1.
- [12] M.F. Ensminger, (1991). *Animals Science.* Inter-state publishers Inc. Denville, Illinious, pp 769-771.
- [13] A.B. Bosch, Poujardieu and R. Rouvier (1992). *Zootechnica and Genetic Potential in Crossbreeding Experiments and Breed Comparisons.* Options Mediterranes-seri seminaries (17): 121-126.
- [14] A. Gorbani, and N.R. Salamatdoust (2011). Comparison of the best breed combination in Iranian Holstein and Brown Swiss crossbred dairy cattle for production traits. *African J. Biotech* 10(85): 19836-19839.
- [15] B.S. Malik and R.P. Singh (2006). Evaluation of Crossbreeding Effects for Wool Traits in Sheep, Asian – *Aus. J. Animal Science* 19 (1); 1536-1540

- [16] O.W. Robinson, B.T. Mac Daniel, and E.J Rincon (1981). Estimation of Direct and Maternal Additive and Heterotic Effects from Crossbreeding Experiments in Animals. *J. Animal Science* 52:44-50.
- [17] Y.M.K. Youssef, M. Baselga, M.H. Khalil, S. Gad-Alla, and M.I. Garcia, (2008). "Evaluation of Litter Traits in a Crossing Project of V-line and Baladi Red Rabbits in Egypt". *Livestock Research for Rural Development* Vol. 20 No. 9. <http://www.Irrd.org/Irrd20/9/yous2013.htm>
- [18] E.A. Afifi, M.H.Khali, A.F. Khadr and Y.M.K. Youssef (2011). "Postweaning Growth Traits and Carcass Performance in Rabbit Crosses", *Journal of Animal Breeding and Genetics*. <http://www.lonlineLibrarywiley.com/doi/10.1111/j.1439-0388.1994.tb00447.x/abstract>
- [19] W.E. Castle, (1922). *Genetic Studies of Rabbits and Rats*. Carnegie Institution of Washington Pub. No. 320.
- [20] M. Baselga, (2004). Genetic Improvement of Mean Rabbit. Programmes and Diffusion. In proc. 8<sup>th</sup> World Rabbit Congress 7-10 September, 2004, Puebla, Mexico, 1-3.
- [21] M. Pascual, and M. Pla, (2007). Changes in Carcass Composition and Meat Quality when Selecting Rabbits for Growth Rate. *Meat Sci.* 77:474-481.
- [22] G.E. Dickerson, (1969). Experimental Approaches in Utilization Breed Resources. *Animal Breed.* Abstr 37:191-202.
- [23] A.I. Ouyed, and J.M. Brun, (2008). "Heterosis, Direct and Maternal Additive Effects on Rabbit Growth and Carcass Characteristics". Proc. 9<sup>th</sup> World Rabbit Congress, Verona, Italy. Pp 195-200.
- [24] Kh.H. El-Bayomi, M.S. El-Tarabary and T.M. Abdel-Hamid, (2012). "Estimation of Heterosis and Combining Ability for some Weaning and Post-weaning Traits in Three Different Breeds of Rabbits". *Journal of American Science*, vol. 8 No. 9, pp 282-288.
- [25] K.M. Elamin and I.A. Yousif (2011). "Evaluation of Litter Traits in Sudanese Rabbits". *Livestock Research for Rural Development*, Vol. 23 No. 9, [www.Irrd.org/irrd23/9/cont2309](http://www.Irrd.org/irrd23/9/cont2309)
- [26] K.A.O. Yamani (1990). "Breeds and Prospects for Research to Improve Rabbit Meat Production in Egypt". *Options Mediterranean's. Sevier seminaries* No. 8 pp 67-73.
- [27] Z.S. Sezendo, (2008). Improving of Reproductive Performance of Rabbit Does in Small and Medium Scale Rabbit Farms. Recommendation for developing Countries. *ACTA Agraria kaps varriensis* (2008) Vol. 12 No. 1, 1-23.
- [28] A.S. Abdel-Azeem, A.M. Abedl-Azim, A.A. Darwish, and E.M. Omar, (2007). "Litter traits in four pure breeds of rabbits and their crosses under prevailing environmental conditions of Egypt". Proceedings of the fifth international conference on Rabbit Production in Hot Climate, Hurghada, Egypt, pp. 39-5 <http://www.fayoum.edu.eg/Agriculture/poultryProduction/Pdf/Abdel-Azim5.Pdf>.
- [29] M.F. Medellin and S.D. Lukefahr (2001). "Bred and heterotic effects on Postweaning traits in atex and New Zealand White Straight Bred and Crossbred Rabbits". *Journal of Animal Science*, Vol. 97, pp. 1173-1178.
- [30] J.M. Brun, and J. Ouhayoun (1989). "Qualities Boucheres de Lapereaux Issues d'un Croisement Dialele de 3 Souches: Interaction du type genetique et de la taille de portee d'origine" *Ann. Zootec*; vol. 43, pp 173-173.
- [31] Y.K. Youssef, M. Baselga, M.H. Khalil, M.E. Omara, and M.L.Garcia, (2009). "Crossbreeding Effects for Post Weaning Growth Traits in a Project of Spanish V-line with Baladi Red Rabbits in Egypt". *Livestock Science*, Vol. 122 No. 2 pp. 302-307.
- [32] A.M. Al-Saef, M.H. Khali, A.H. Al-Homidan, S.N. Al-Dobaib, K.A. Al-Sobayil, M.I. Garcia and M. Baselga, (2008). "Cross Breeding Effects for Litter and Lactation Traits in Saudi Project to Develop New Lines of Rabbits Suitable for Hot Climates". *Livestock Science*, Vol. 118 No. 3 pp 238-246.
- [33] E.J. Eisen and B.S. Durant (1980). Effect of Parity on Reproductive Performance of Mice Selected for Litter Size on Body Weight. *Res. Brasil-Genet.* III, I, 1-19 (Brazil J. Genetics).