

Fighting against Desertification: Studying Physiological Markers of Halophytes Response to Salinity

AMROUNE Mohamed Lamine*, BELKHODJA Moulay
Laboratory of Plant Physiology, University of Oran 1 Ahmed Benbella
*Email: amrounemohamed2003@yahoo.fr

Abstract – This work is a comparative study of photosynthetic gaseous exchange and respiratory activity of isolated leaf chloroplasts and leaf discs on a halophytic plant (*Atriplex halimus* L.). Plants were stressed with a saline solution consisting of 400 meq.l⁻¹ NaCl+CaCl₂ and 600 meq.l⁻¹ Hoagland nutrient solution added to diluted (50%) and non-diluted seawater. Gaseous exchange was measured during 30 min using Warburg device. Oxygen released throughout photosynthetic reaction in daylight conditions and oxygen absorbed throughout respiratory activity in darkness conditions was recorded.

Results revealed that seawater (or NaCl+CaCl₂) salinity affects respiratory and photosynthetic activities. They however vary according to medium concentration and stress type.

Atriplex halimus L. leaf chloroplasts gaseous exchange shows indeed an oxygen absorption decrease with a 400 meq.l⁻¹ NaCl+CaCl₂ concentration. Watering with non-diluted seawater causes respiratory activity slow down on *Atriplex halimus* L. leaf chloroplasts.

Atriplex halimus L. leaf chloroplasts photosynthetic activity is more intense under 400 meq.l⁻¹ NaCl+CaCl₂ stress.

Keywords – *Atriplex halimus* L., Gaseous Exchange, Leaf Chloroplasts, Leaf Discs, NaCl+CaCl₂, Oxygen Breathing, Oxygen Photosynthesis, Salinity, Seawater, Warburg Device.

I. INTRODUCTION

After a long lasting drought, Algerian soil has become halomorphic. This characteristic is specific to arid and semi arid countries [Osman, 1982; Munns, 2002]. On Earth, salinity covers about 954.8 millions hectares and 27% of them are farmland [Hamdy *et al.*, 1999]. The Mediterranean Basin is affected by 16 millions hectares of salted land: 3.2 millions are in Algeria [Hamdy, 1999]. In these ecosystems, salinity has become a limiting factor for crop development [Tal, 1984; Hamdy *et al.*, 2002]. This is due to abnormal high soluble salt rate [Poljakof-Mayber & Gale, 1975; Kolahchi & Jalali, 2007]. Actually, salt excess induces nutritional poisoning phenomena due to absorption disorders [Zid, 1986; Meloni, 2001]. It is also due to accumulated compounds becoming toxic and disrupting plant metabolism [Sanchez Gonzalez & Agoyo, 1992; Liu & Zhu, 1998] and due to high osmotic potential [Immamulhaq *et al.*, 1984; Yokoi *et al.*, 2002; Moinuddin *et al.*, 2005]. This changing biotope has led to species extinction [Chamard, 1993; Pitman & Lauchli, 2002] while others have acquired resistance mechanisms [Christian, 1992; Edward *et al.*, 1999]. In addition, soils structural fertility and stability have been modified [Valentin, 1994; Benkhelifa, 1997; Reguieg, 2007].

In practice, restoring salt damaged soils requires many actions. One of them is introducing legumes and

halophytes. In weak rainfall regions, this eco-physiological approach allows recovering soil fertility [Sing *et al.*, 1982; Belkhodja, 2004] through symbiotic nitrogen fixation by legumes [Osman *et al.*, 1986; Jensen, 1986; Belkhodja *et al.*, 2002], and through desalinization and soil fixing by halophytes [Ashraf *et al.*, 1987; Le Houerou, 1992; Gorham, 1996; Batanouny, 1996]. Choosing legumes as broad beans and halophytes as *Atriplex halimus* L. is based on every species several advantages. In addition to advantages already mentioned, they also are a source of animal protein [Polignano *et al.*, 1991] and are tolerant to salinity, especially *Atriplex* [Le Houerou, 1991]. Moreover, *Atriplex* species seem nowadays to be the most adapted to steady and increase forage production in arid and semi-arid climate [Froment, 1972; Kinet *et al.*, 1998].

Introducing these species in saline ecosystems requires understanding certain mechanisms such as physiological mechanisms involved in plant resistance to abiotic constraint. In accordance with [Boucaud *et al.*, 1978; Girija, 2002], plant behavior varies indeed according to surrounding environment salinity: whether steady, increasing continuously or fluctuating.

II. MATERIAL AND METHODS

Plant material

The seeds of *Atriplex*, *Atriplex halimus* L., used in our experiments is taken starting from shrubs pushing in the campus of the University of Senia Oran (Algeria).

Atriplex seeds were sown in cells. Then seedlings were carefully transferred into flowerpots with sand substrate (2/3) and industrial peat (1/3). Plants were monitored during 157 days watered with 1/1000 diluted Hoagland (1938) nutrient solution before receiving saline treatment.

Experimental process

After 157 days, 400 meq.l⁻¹ and 600 meq.l⁻¹ NaCl+CaCl₂ saline solutions were poured over *Atriplex*. After one-week stress, leaves were carefully removed to extract chloroplasts and leaf discs. With 1 ml chloroplasts suspension and 10 leaf discs, gaseous exchange was monitored through Warburg device. Photosynthetic oxygen was recorded in daylight while respiratory oxygen was recorded in darkness conditions, both during 30 min.

Extraction of the chloroplasts

This operation proceeded according to the process of Walker (1970). This process normally makes it possible to obtain a suspension of chloroplasts of class I (complete), able to fix in the proportion from 50 to 80%, a CO₂ high rate.

For that, preparations took place:

Medium of crushing

Sorbitol 0.33 M
MgCl₂ 0.005 M
Na₄P₂O₇, 10:2 O 0.01 M

Brought the pH of the mixture above to 6.5 with HCl, and to add, right before employment, of the sodium isoascorbate until a concentration 0.002 Mr.

Medium of handing-over in suspension

Sorbitol 0.33 M
MgCl₂ 0.001 M
MnCl₂ 0.001 M
EDTA 0.002 M

HEPES (tampon acid hydroxyethylpiperazine-ethane sulfonic) 0.05 Mr.

Brought the pH to 7.6 with soda.

Extraction itself

For the extractions of the chloroplasts, the sheets are taken median part of each treated plant. 8 G of sheets are crushed in a crusher with 100 ml of medium of crushing. Then, the medium obtained is dried through 2 layers of gauze in a beaker and is filtered through 8 layers of gauze using a vacuum pump; the recovered solution is versed in a test tube then last in a blender during 3 to 5 seconds. This medium is versed in two tubes with centrifugation with equal volumes. The whole is centrifuged at low temperature (0° C) at a speed of 4000 tours/min during 90 seconds; in each tube is added 1 ml of medium of handing-over in suspension to the base recovered using a pipette (1 ml). Lastly, the suspension of chloroplasts is taken again using a syringe in two tubes of Eppendorf to then put them in a refrigerator while waiting for measurements of the gaseous exchange.

III. RESULTS


Variations in respiratory and photosynthetic gaseous exchange
I- CHLOROPLASTS GASEOUS EXCHANGE MEASURED DURING 30 MIN.
A. Respiratory gaseous exchange


Fig. 1. Variation of respiratory oxygen XRO₂ (mm³) absorbed during 30 min by 1 ml chloroplasts suspension extracted from 165 days old *Atriplex halimus* L. leaves stressed with NaCl+CaCl₂.

Fig. 1 shows that the amount of oxygen absorbed by control plants chloroplasts increases to a maximum of 46.248 mm³ during the first five minutes. Then the amount decreases to 8.742 mm³ during the next five minutes. Oxygen absorption fall continues for 20 minutes gaseous exchange until the end of experiment.

With chloroplasts from plants fed with 400 meq.l⁻¹ NaCl+CaCl₂, the amount of respiratory oxygen increases to a maximum of 14.865 mm³ during the first five minutes and then decreases to -9.644 mm³ during the last 20 minutes. With chloroplasts from plants fed with 600 meq.l⁻¹ NaCl+CaCl₂, the amount of oxygen exchange increases to a maximum of 20.899 mm³ after five minutes and then decreases until the end of experiment.


B. Photosynthetic gaseous exchange


Fig. 2. Variation of photosynthetic oxygen XPO₂ (mm³) released during 30 min by 1 ml chloroplasts suspension extracted from 165 days old *Atriplex halimus* L. leaves stressed with NaCl+CaCl₂.

Fig. 2 shows that the amounts of photosynthetic oxygen released by control plants and stressed plants have a parallel evolution after 10 minutes of gaseous exchange. Control plants chloroplasts release goes up to 27.05 mm³ photosynthetic oxygen during 15 minutes and then falls down to 8.32 mm³.

In stress conditions, oxygen released is kept on a high level on chloroplasts from plants watered with 400 meq.l⁻¹ NaCl+CaCl₂ concentration and then decreases slowly after 20 minutes.

In 600 meq.l⁻¹ NaCl+CaCl₂ medium, photosynthetic activity evolves in a intermediary way between chloroplasts from control plants and from 600 meq.l⁻¹ NaCl+CaCl₂ stressed plants. It should be noted that for stressed plants, amount of oxygen released has a similar evolution than amount recorded with 400 meq.l⁻¹ salt treated plants.

II- LEAF DISCS GASEOUS EXCHANGE MEASURED DURING 30 MIN.
A. Photosynthetic gaseous exchange

Fig. 3 shows variations in oxygen released during photosynthetic reaction in daylight conditions. In the beginning of the experiment, oxygen released by leaf discs increases with all plants, whatever the treatment has been. After 5 minutes, a maximum of oxygen is released by leaf discs recording 34.966 mm³ with control plants, 21.465 mm³ with 400 meq.l⁻¹ NaCl+CaCl₂ treated plants and 30.628 mm³ with 600 meq.l⁻¹ NaCl+CaCl₂ treated plants.




Fig.3. Variation of photosynthetic oxygen XPO₂ (mm³) released during 30 min by 165 days old *Atriplex halimus* L. leaf discs stressed with NaCl+CaCl₂.

After reaching these values, released photosynthetic oxygen decreases slowly for all plants during 10 minutes. Then, oxygen production increases back very rapidly with control plants leaf discs, reaching 70.387 mm³ after 30 minutes. On the other hand, oxygen released becomes steady with leaf discs from 400 meq.l⁻¹ NaCl+CaCl₂ treated plants while it slows down with leaf discs from 600 meq.l⁻¹ NaCl+CaCl₂ watered plants.

B. Respiratory gaseous exchange

Fig. 4 shows variations in oxygen absorbed by leaf discs during respiratory activity in darkness conditions. After 5 minutes, control plants leaf discs absorbed a maximum of -39.806 mm³, 400 meq.l⁻¹ NaCl+CaCl₂ stressed plants leaf discs absorbed a maximum of -22.96 mm³ and plants fed with 600 meq.l⁻¹ NaCl+CaCl₂ leaf discs a maximum of -31.82 mm³. Beyond this point, oxygen absorbed by control plants leaf discs slowed down during 15 minutes to reach -7.084 mm³ at the end of the experiment.




Fig.4. Variation of respiratory oxygen XRO₂ (mm³) absorbed during 30 min by 165 days old *Atriplex halimus* L. leaf discs stressed with NaCl+CaCl₂.

Respiratory activity of 400 meq.l⁻¹ NaCl+CaCl₂ stressed plants leaf discs slowed down until the end of the experiment reaching a -4.294 mm³ value. Same phenomenon was recorded with leaf discs from plants fed with 600 meq.l⁻¹ NaCl+CaCl₂.

IV. CONCLUSION AND GENERAL DISCUSSION

The purpose of this work was to monitor and compare photosynthetic and respiratory gaseous exchange of a

halophytic plant (*Atriplex*) leaf discs and isolated chloroplasts. Plants were cultivated in flowerpots stored in a greenhouse to be watered with 400 meq.l⁻¹ saline solution and 600 meq.l⁻¹ NaCl+CaCl₂ gaseous exchange activity was measured through Warburg device during 30 minutes.

Results recorded assess that *Atriplex halimus* L. chloroplasts variations in oxygen absorption during respiratory activity and in oxygen release during photosynthetic activity are due to presence of salt with different concentrations in nutrient solution.

• *Atriplex chloroplasts gaseous exchange*

* concerning respiratory gaseous exchange

While plants are stressed with NaCl+CaCl₂, respiratory oxygen absorbed by chloroplasts measured in darkness conditions during 30 minutes shows weak respiratory activity during the first 10 minutes compared to control plants absorbing up to 46.248 mm³ oxygen. But during the last 20 minutes respiratory activity decreases very slowly until the end of the experiment whatever the salt concentration is.

* concerning photosynthetic gaseous exchange

Photosynthetic activity goes up to a maximum after 20 minutes whatever the salt concentration is: 38.968 mm³ with 400 meq.l⁻¹ NaCl+CaCl₂ treated plants and 27.28 mm³ with 600 meq.l⁻¹ NaCl+CaCl₂ treated plants. Control plants chloroplasts release a maximum of oxygen after 25 minutes (27.05 mm³).

• *Atriplex leaf discs gaseous exchange*

* concerning photosynthetic gaseous exchange

Control plants leaf discs show a high photosynthetic activity during the whole experiment (70.387 mm³). Salinity decreases leaf discs oxygen absorption on plants watered with 400 meq.l⁻¹ and 600 meq.l⁻¹ NaCl+CaCl₂ (21.465 mm³ and 30.628 mm³ respectively).

* concerning respiratory gaseous exchange

Leaf discs respiratory activity of NaCl+CaCl₂ stressed plants is weak (-22.96 mm³ with 400 meq.l⁻¹ and -31.82 mm³ with 600 meq.l⁻¹) compared to control plants (-39.806 mm³).

Gaseous exchange recorded after 30 minutes allows to compare respiratory and photosynthetic activities:

- *Atriplex* chloroplasts respiratory gaseous absorption decreases as salt concentration increases.
- *Atriplex* chloroplasts photosynthetic oxygen release increases as salt concentration equals to 400 meq.l⁻¹.

Respiratory and photosynthetic gaseous exchange of *Atriplex halimus* L. chloroplasts from plants watered with saline solution has been the purpose of a certain number of laboratory works. Conclusions of works conducted on leaf discs assess that salinity has an impact on photosynthetic and respiratory gaseous exchange [Amroune, 2000]. With similar experimental conditions, it has been assessed that leaf discs absorb more oxygen during respiratory activity when plants are stressed with intermediate salinity (i.e. 400 meq.l⁻¹ NaCl+CaCl₂). [Kabou, 1997] and [Abderrahmane, 2001] have showed that respiratory activity recorded with *Atriplex halimus* L. leaf discs increases as salt concentration increases. Moreover, these works report that photosynthetic oxygen release is very

important under high salt concentration stressing conditions. Respiratory and photosynthetic exchange negative values recorded in our works likely reflect photorespiration phenomenon, a process occurring during daylight and characterized by oxygen absorption and CO₂ release.

[Bensafi & Bouchenak, 1997] have also established that salinity causes respiratory gaseous exchange increase with broad beans.

[Chahrouf, 2002] has concluded that salinity has an impact on *Atriplex halimus* L. chloroplasts respiratory and photosynthetic activities. He has also concluded that amount of oxygen absorbed during respiratory activity and released during photosynthesis vary throughout duration according to salt concentration, stress type and ecotype. [Kheloufi, 2004] has demonstrated that broad beans leaf chloroplasts photosynthesis reaches higher values compared to control plants when fed with 400 meq.l⁻¹ NaCl+CaCl₂. No photosynthetic activity was recorded when salt concentration equals to 600 meq.l⁻¹ NaCl+CaCl₂.

[Björkman & Berry, 1973] have compared the impact of several factors on two *Atriplex* species photosynthesis: *Atriplex patula* (with a C₃ photosynthetic metabolism type) and *Atriplex rosea* (with a C₄ photosynthetic metabolism type). Both researchers have reported that C₄ plants are more efficient to photosynthesis under heavy sunlight and high temperature than C₃ plants. Moreover, C₄ metabolism plants continue photosynthesizing while CO₂ concentration within intercellular spaces is very low, which is the case when stomata are closed.

According to results, *Atriplex halimus* leaf chloroplasts respiratory activity decreases because of salinity. This may reflect an adaptation mechanism to stress, one of halophytic plants characteristics.

Atriplex halimus L. leaf chloroplasts photosynthetic activity is very intense whatever the salt concentration is.

After comparing isolated chloroplasts and leaf discs gaseous exchange, results assess that *Atriplex halimus* leaf discs gaseous exchange is more active, probably due to the photorespiration effect taking place at the same time than photosynthesis. The phenomenon requires chloroplasts, mitochondria and peroxisomes functions. Photorespiration phenomenon takes place in case of plants being in stress conditions such as drought and salinity. This is to produce a maximum of energy and to compensate energy loss due to photosynthetic intensity decrease. [Costa *et al.*, 1978] have notified that photorespiration increases as oxygen partial concentration does.

REFERENCES

- [1] Abderrahmane, L. 2001. Action de la salinité sur les échanges gazeux photosynthétiques et respiratoires chez l'*Atriplex halimus* au stade juvénile, Mémoire de D.E.S. Université d'Oran, 71 p.
- [2] Amroune, M A. 2000. Action de la salinité sur les échanges gazeux respiratoires et photosynthétiques chez l'*Atriplex halimus* L. au stade juvénile de la plante, Mémoire de D.E.S. Université d'Oran, 65 p.
- [3] Ashraf, M. 1987. McNEILLY T and BRADSHAW. Selection and heritability of tolerance of NaCl in four forage species. *Sci.*, 227, p. 232-234.
- [4] Batanouny, K H. 1996. Ecophysiology of Halophytes and Their Use in the Arab World. Halophytes and Biosaline Agriculture. Choukr-Allah, R.; Hamdy, and Malcolm, C.V, p.73-91.
- [5] Belkhdja, M. 2004. The physiological response of faba bean (*Vicia faba* L.) to the salinity: a study of the plant transpiration. *Egyptian J. of Applied Sci.*, vol. 5, N° 10, 8p.
- [6] Belkhdja, M., R. Chadli, and Z. Hadj Ziane. 2002. The ionic response of faba bean (*Vicia faba* L.) to the salt stress. *Egyptian J. of Applied Sciences*, N° 15, (12), 422-437.
- [7] Benkhelifa, M. 1997. Influence de la bentonite sur les caractéristiques physiques, hydriques et mécaniques d'un sol sableux du plateau de Mostaganem. Thèse de Magister INA d'EL Harrach. Alger, 114p.
- [8] Bensafi, N., et F. Bouchenak. 1997. Influence de la salinité sur la chlorophylle et la quantité d'oxygène échangée des disques foliaires de la Fève, Mémoire de D.E.S. Université d'Oran, 5p.
- [9] Björkman, O., and J. Berry. 1973. High efficiency photosynthesis. *Scientific American* 229, 80-90.
- [10] Boucaud, J., et Irwin Ungara. 1978. Halophilie et résistance au sel dans le genre *Suaeda* forsk. *Soc. Bot. Fr. Act. Bot.* 34, 23-35.
- [11] Chahrouf, F. 2002. Les échanges gazeux photosynthétiques et respiratoires des chloroplastes foliaires de l'*Atriplex halimus* L. stressée à la salinité, Thèse de Magister. Univer. d'Oran, 100 p.
- [12] Chamard, P. 1993. Environnement et développement. Références particulières aux états sahéliens membres du CILSS. *Re. Sécheresse*, 4, p. 1723-1728.
- [13] Chretien, D. 1992. La résistance au sel chez le joboba. (*Simmondsia chinensis* Linck, *Schneider*): Croissance et modification du contenu lipoprotéique de cal cultivés en présence d'une teneur élevée en NaCl. Thèse Dr. Univer. Pierre et Marie Curie. Paris VI. UFR des sciences de la vie, 116 p.
- [14] Coste, C. 1978. Photosynthèse et production végétale, Gauthier-Villars, BORDAS, Paris, 0219 780 109, ISBN: 2-04-010 194-2.
- [15] Edward, P., E. P. Glenn, J. J. Brown, and E. J. Blumwald. 1999. Salt tolerance and Crop Potential of Halophytes Critical Reviews in Plant Sciences Volume 18, Issue 2, March 1999, 227-255.
- [16] Froment, D. 1972. Etablissement des cultures fourragères d'*Atriplex* en Tunisie centrale. *Bull. Recherche, Agro. C.M.E.L.O.N. Vol. Extrac. Ed. À l'occasion de la semence d'étude de prod. Méditerranée*, pp: 590-600.
- [17] Jensen, E. S. 1986. Symbiotic N₂-Fixation in pea and field bean estimated by N₁₅ fertilizer dilution in field experiment with burley as a reference crop. *Plant and soil*, 92, p 3-13.
- [18] Hamdy, A., H. Lieth, and Z. Mezher. 1999. Halophyte performance under high salinity levels: an overview. Saline irrigation: halophyte production and utilisation; 20-58.
- [19] Hamdy, A. 1999. Saline irrigation assessment and management for a sustainable use saline irrigation: halophyte production and utilisation; 152-226.
- [20] Hamdy, A., N. Katerji, J. W. Van Hoorn, and M. Mastrorilli. 2002. Mediterranean crop responses to water and soil salinity, Ecophysiological and agronomic analyses; Options Méditerranéennes Série B 36, p. 1-3.
- [21] Immamulhaq, S. M., and F. Larher. 1984. Effect of maintaining a constant Na: Ca ratio on the growth, ion balance and organic solute status of NaCl treated cowpea (*Vigna sinensis* L.). *Z. Pflanzenphys.* 113, 163-176.
- [22] Giriya, C., B N. Smith, P M. Swamy. 2002. Interactive effects of sodium chloride and calcium chloride on the accumulation of praline and glycinebetaine in peanut (*Arachis hypogaea* L.). *Environ. Exp. Bot.* 47:1-10.
- [23] Gorham, J. 1996. Mechanisms of Salt-tolerance of Halophytes. Halophytes and Biosaline Agriculture. Choukr-Allah, R; Hamdy, and Malcolm, C.V, p.31-50.
- [24] Kabou, D. 1997. Influence de la chlorure de sodium et calcium (NaCl+CaCl₂) sur les échanges gazeux photosynthétiques chez quelques lignées de fève, Mémoire de D.E.S, université d'Es-Sénia, 56 p.
- [25] Kheloufi, A. 2004. Caractérisation physiologique de la résistance de la fève (*Vicia faba* var. *Aguadulce*) à la salinité, Mémoire de DES, Univ. d'Oran.
- [26] Kinet, J M., F. Benrebaha, Bouzid, S. Lailhacac et P. Dutuit. 1998. Réseau Atriplex. Atelier Biotechnologie et Ecologie pour une sécurité alimentaire accrue en régions arides et semi arides. *Cahiers de l'Agriculture*, 7, p. 53-62.

- [27] Kolahchi, Z., and M. Jalali. 2007. Effect of water quality on the leaching of potassium from sandy soil. *Journal of Arid Environments* Volume 68, Issue 4, 624-639.
- [28] Le houero, H N. 1991. The role of salt bushes (*Atriplex* spp.) in arid land rehabilitation in the Mediterranean Basin: a review *Agroforestry Systems*, 18. 107-148.
- [29] Le houero, H N. 1992. The role of salt bushes (*Atriplex* spp.) in arid land rehabilitation in the Mediterranean basin: A review *Agroforestry Systems*, 18, p. 107-148.
- [30] Liu, J., and J K Zhu. 1998. A calcium sensor homolog required for plant salt tolerance. *Science*, Vol. 280. no. 5371, pp. 1943-1945.
- [31] Meloni, DA., MA. Oliva, H A. Ruiz, C A. Martinez. 2001. Contribution of proline and inorganic solutes to osmotic adjustment in cotton under salt stress. *J. Plant Nutr.* 24:599-612.
- [32] Moinuddin, A., R A. Fischer, K D. Sayre, and M P. Reynolds. 2005. Osmotic Adjustment in Wheat in Relation to Grain Yield under Water Deficit Environments. *Agro. J.* 97:1062-1071.
- [33] Munns, R. 2002. Comparative physiology of salt and water stress. *Plant Cell Environ*; 25: 239-250.
- [34] Osman, A. 1982. The future of arid zones. 12th congress Int. Sci. of Soil. Managing and Resources; 101-105.
- [35] Osman, A K., H E. Habib Ibrahim, and MC. Saxena. 1986. Seasonal pattern of symbiotic Nitrogen fixation in *Faba beans* and it's Association with some physiological traits. *Fabis Newsletters*, n° 14.
- [36] Pitman, M G., and A. Läuchli. 2002. Global impacts of salinity and agricultural ecosystem *In* A Läuchli, U Lüttge eds, *Salinity: Environment-Plants-Molecules*. Kluwer Academic, Dordrecht, The Netherlands, pp 3-20.
- [37] Poljakof, M., and Gale. 1975. Morphological and anatomical changes in plants as a response to salinity stress. In "Plants and in saline environments". *Ecological studies*, 15, Springer Verlag, N-Y, B, 95-117.
- [38] Polignano, G B., R. Splendido, and P. Ugenti. 1991. Protein polymorphism among genotypes of Faba bean (*Vicia faba* L.) from Ethiopia and Afghanistan. *Fabis*, p. 8-11.
- [39] Reguieg Yssad, H A. 2007. Effet de la bentonite sur les sols sableux de la région de Mostaganem, étude de comportement d'une association blé pois chiche. Thèse de doctorat INA EL Harrach. Alger, p 94-100.
- [40] Sanchez, A., and AL. Gonzalez Utor. 1992. Quantitative determination of changes induced by NaCl in vacuoles and cellular size of *Lycopersicum esculentum* root cells. *Plant Cel. and Env.*, 15, p. 867-870.
- [41] Sing, CB., R. Ramgiris, and SB. Singh. 1982. Faba bean a high protein potential pulse crop Indian farming. *Fabis Newsletter*, 21, p 26.
- [42] Tal, M. 1984. Selection for stress tolerance. In hand book of *Plant Cell Culture*; Techniques for propagation and breeding (D.A. Evans, W.R. sharp, P.V. Ammirato and Y. Yamadou, eds). Vol. I. p. 461-488. Mac Millan publishing co. New York. ISBN0-02-949230-0.
- [43] Valentin, C. 1994. Sécheresse et érosion au Sahel. *Rev. Sècheresse*, 5, p 191- 98.
- [44] Yokoi, S., RA. Bressan, and PM. Hasegawa. 2002. Salt stress tolerance of plants. *JIRCAS Working Report*, 25-33.
- [45] Zid, E., et C. Grignon. 1986. Tolérance du bigaradier à la salinité. *Acta Oecologica Planta.*, 7, 21p. 407-416.