

Effect of Tomato *Lycopersicon Esculentum* (MILL) Varieties on Development Time, Fecundity and Longevity of Red Spider Mite *Tetranychus Evansi* (Baker and Pritchard)

MUSAH S. M.Ministry of Agriculture, Livestock and
Fisheries, Molo Sub- County, Kenya.

Author for correspondence

(Email ID : msmatika@yahoo.com)

KAMAU A. W.Department of Crop, Horticulture and Soil
Sciences, Egerton University, Njoro,
Kenya**MUNENE M**Kenya Agricultural Research Institute,
Njoro, Kenya

Abstract – Tomato *Lycopersicon esculentum* (Mill), is an important source of income to many small scale farmers in Kenya. However, the production is constrained by several pests with Red spider mite *Tetranychus evansi* (Baker & Pritchard) being the most important dry season pest in Eastern and Southern Africa. Control of *T. evansi* has mainly been by chemicals sprays, which has resulted in problem of pest resistance and pollution of the environment. This study was therefore an attempt to look into alternative environmentally friendly ways of controlling *T. evansi* in tomatoes. To determine the effect of tomato varieties on development time, fecundity and longevity of *T. evansi*, an experiment was conducted in a conviron incubator maintained at 25 ± 2 °C ,50-70 % relative humidity and photoperiod of 12:12 L:D in a complete randomized design (CRD) arrangement replicated four times in Kenya Agricultural Research Institute (KARI), Njoro, Kenya. The results showed that the, mortality rate for protonymph, deutonymph and adult stages was highest in Wild type (95%) and lowest in variety Money maker (50%). In addition, the average number of eggs per female mite was significantly higher on variety Money maker (52.4) and Riogrande (48.6), while the least number of eggs was recorded on Wild type (9.1). The results of this study indicate that antibiotic effect existed especially in Wild type. This shows that there is potential for host plant resistance for the control of *T. evansi* on tomatoes and may form part of an IPM programme.

Keywords – Duetonymph, Host Plant Resistance, IPM, *Lycopersicon Esculentum*, Protonymph, *Tetranychus Evansi*.

I. INTRODUCTION

Tomato *Lycopersicon esculentum* (Mill) is the world's most popular vegetable, with an annual world production of 80 million metric tons (FAO, 2001). Kenya produces an estimated 318,639 metric tons and earning cash value of Ksh 5.1 billion (KARI, 1996). In Kenya, it is one of the most important local vegetable crops ranking second to *Brassica* (cabbage and kales) in quantities produced and value (KARI, 1996). The increased popularity in tomato production may be attributed to both its high yield potential, high prices and a continuous source of income (Kamau, 1985), and perhaps more than any other vegetable, it has a higher monetary return per unit area. Tomato growing is thus an important undertaking by low to moderate class earners and goes along way in supporting the fight against poverty and nutritional improvement in Kenya.

In Kenya, tomato is grown in almost all arable areas including semi arid and arid areas using irrigation. However, the major tomato production constraints in Kenya are Red spider mite (*T. evansi*), bacterial wilt, late blight, plant parasitic nematodes, insect pests, lack of high yielding varieties and poor agronomic practices (KARI, 1996).

Red spider mite (*T. evansi*), is a major pest of solanaceous crops including tomato, pepper, eggplant, tobacco and nightshade (Quereshi, *et al.*, 1969., Ramalho and Flechtmann, 1979). It is the most important dry season pest of tomatoes in eastern and southern Africa. Kamau, 1985, reported that these are serious pests of tomato grown in areas where the crop experience periods of hot and warm dry weather. *Tetranychus evansi*. causes serious damage to the tomatoes by reducing their yields and affecting quality, not to mention the cost of pesticides used by the farmers (Saunyama and Knapp, 2003). Compounding their effects is the fact that in many instances, farmers have listed Red spider mites as a disease (Yang *et al.*, 2004) because they are unable to recognize their minute sizes on crop surfaces. Resistant tomato varieties to Red spider mites have been reported in some countries. For instance, in USA, Kalohi lines recorded high tolerance to mite injury (Varela *et al.*, 2003). In Brazil, some promising tomato genotypes tolerant to *T. evansi* have been identified (Varela *et al.*, 2003). There are many tomato varieties commercially available to growers in Kenya; however these varieties have not been screened to compare their resistance levels to Red spider mites. This study was conducted to determine the effect of tomato varieties on *T. evansi* development time, fecundity and longevity.

II. MATERIALS AND METHODS

Mite Culture

A stock culture of Red spider mite *T. evansi* was established in green house at KARI- Njoro ($0^{\circ} 20'S$ $35^{\circ} 56'E$, 2164 m above sea level). The initial culture was collected from Horticulture Department (Field 3), Egerton University- Njoro, Kenya and reared on potato plants. The reared red spider mites were later used in laboratory experiment carried out at KARI- Njoro. Red spider mite, *T. evansi* was identified in consultation with International

Centre of Insect Physiology and Ecology (ICIPE) – Nairobi.

Development Time From Egg to Adult

Studies on the effect of tomato varieties on the development time from egg to adult of *T. evansi* were done in a conviron incubator maintained at $25 \pm 2^\circ\text{C}$, 50-70 % relative humidity and photoperiod 12:12 L:D (Helle and Sabelis, 1985). Small Petri dishes (100 mm diameter) were loaded half with cotton wool and kept wet by adding 5ml of clean water in the morning and evening. Ten leaf discs from each tomato variety with a diameter of 20 mm each were punched from the tomato varieties using a cork borer. The leaf discs were placed in individual Petri dishes, the underside facing up. Thin layer of vaseline was placed on the edges of the leaf disc to prevent the mite from walking a way from the leaf disc and drowning. Ten female mites were taken from the colony, put on each leaf disc for each tomato variety separately and left for six hours to allow egg laying. Afterwards, eggs were removed (killed) using fine camel hair brush leaving only two eggs per leaf disc. Petri dishes containing leaf discs were then placed in an incubator in a complete randomized design (CRD) arrangement replicated four times. The leaf discs were checked after every 12 hours with the aid of a dissecting microscope and development time from egg to larva, protonymph, deutonymph and adult, as well as adult survival and sex were noted and recorded. Leaf discs were changed after every three days to provide fresh leaves for mites to feed on.

Fecundity and Longevity

Studies on the effect of tomato varieties on fecundity and longevity of *T. evansi* were done in a conviron incubator maintained at $25 \pm 2^\circ\text{C}$, 50-70 % relative humidity and photoperiod 12:12 L:D (Helle and Sabelis, 1985). Small Petri dishes (100mm diameter) were loaded half with cotton wool and kept wet by adding 5 ml of clean water in the morning and evening. Ten leaf discs with a diameter of 20 mm from each tomato variety were punched using a cork borer. The leaf discs were placed in individual petri dishes, the underside facing up. Thin layer of vaseline was placed on the edges of the leaf disc to prevent the mite from walking a way from the leaf disc and drowning. One female deutonymph was placed on each leaf disc for each of the ten leaf disc per tomato variety. Petri dishes containing leaf discs were then placed in an incubator in a complete randomized design (CRD) arrangement replicated four times. The number of eggs was counted and recorded using dissecting microscope after every 48 hours and killed until the mite died. The fecundity and adult longevity on each leaf disc was recorded. Leaf discs were changed after every three days to provide fresh leaves for mites to feed on.

Data Analysis

Data collected was organized and Analysis of variance (ANOVA) was done using the General linear model (GLM) procedure of SAS 2001 version 8.02. Means were separated using Tukey's Studentized Range Test.

III. RESULTS

There was no significant difference ($P>0.05$) on incubation period, and duration of larval stage in all the varieties tested (Table 1a). However, there was significant difference ($P<0.05$) on duration of protonymph, and duetonymph stages and as well as adult survival among tomato varieties (Table 1a and Table 1b). The mortality rate of protonymph, duetonymph and adult stages was highest in wild type with 44.4%, 70.0% and 66.7% respectively and lowest in Money maker with 15.0%, 23.5% and 23.1% respectively.

The total development period (egg to adult) ranged between 8.9 to 11.0 days (Table 1a and 1b). The average number of eggs laid per female mite was significantly higher on Money maker which was not significantly different ($P>0.05$) from Riogrande (Table 2). The least number of eggs was recorded on Wild variety which was significantly different from all the varieties.

IV. DISCUSSIONS

The results on incubation period and duration of larval stage may be because the mites are not feeding on the host plant and therefore there is possibly that there is neither morphological nor chemical effect on the mite development at this stage. The significance difference in the duration of time taken on these stages (Protonymph, Duetonymph and adult) may be because at these stages the mite has started feeding on the host plant. The difference in duration may therefore be attributed to the different morphological traits such as trichomes and surface waxes as well as secondary plant metabolites on different tomato varieties (Anathakrishnan, 2001). The difference in mortality rate in the present study may be due to difference in host plant characteristics. This is similar to what was reported by Chatzivasileiadis and Sabelis, 1997,1998., Chatzivasileiadis *et al.*, 1999., Maluf *et al.*, 2001., Cedola and Sanchez, 2003) where compounds from tomato trichome were reported to produce exudates such as methyl ketons and zingiberene that were reported to be toxic to spider mites. High densities of glandular type IV trichomes on *L. hirsutum*, and non-glandular type V trichomes on *L. esculentum* were also found by Simmons and Gurr (2005). Glandular trichome types I, IV, VI and VII are known to release secretions on touch that are sticky and toxic (Van haren *et al.*, 1987). Nihoul, 1993, Kennedy, 2003, Simmons and Gurr, 2005). The secretions immobilize mites or kill them.

The total development period (egg to adult) ranged between 8.9 to 11.0 days. Similar results were reported by Wesonga *et al.*, (2005). It was noted that the period of *T. evansi* from egg to adult on tomato leaves ranged from 6.5 to 11.5 days. Meyer, (1996) also reported that the life cycle is 9 to 12 days resulting to 24-30 generations in a year. The difference in time taken for eggs to hatch and larva that completed the cycle indicated that development and population increase of *T. evansi* depends on host type. In this case Money Maker is a more suitable host compared to Wild type and other commercial varieties

tested because 50% of the larvae reached the adult stage. Similar results were also reported by Isutsa *et al.*, (2006) where it was reported that 58 % of the larvae on Money maker reached adult stage. Host plants can exert profound effects on the biology of spider mites including *T. evansi* Jeppson *et al.*, (1975). In this study the average ratio of male to female was 1:2.6. This is less compared to what was reported by Qureshi *et al.* (1969) that the ratio of male to female was about 1:10. These difference may be due to difference in test plant varieties used and also different laboratory conditions i.e. temperature and relative humidity. However, the results seems to be almost similar to work done by Moutia (1958) who reported that the sex ratio on tomato plants in the field was 1:3.3. The short life cycle and high ratio of female to male indicate that on suitable host and favorable climatic conditions *T. evansi* can reproduce rapidly and cause severe crop losses.

In the present study, highest mortality was recorded on Wild type while lowest mortality was recorded on variety Money maker. Similar results were also reported by Snyder and Carter (1984 and 1985) where it was reported that mite survival was lower in *L. hirsutum* var. *hirsutum*

that possessed fewer type VI trichomes than *L. esculentum*, and concluded that trichome type VI could be responsible for the high levels of resistance in this species. The results on fecundity and longevity of *T. evansi* agree with what was reported by Isutsa *et al.*, (2006) where the highest total number of eggs laid by an individual female mite was reported on Money maker. From the results, Money maker seems to be a preferred host plant for egg laying as compared to wild type. Similar results were also reported by Murungi *et al.*, 2009 where preference to Money maker was attributed to trichome density and plant emitted volatiles. This may be attributed to the different morphological traits such as trichomes and surface waxes as well as secondary plant metabolites (Anathakrishnan, 2001). The wild type may also be releasing some chemicals that repel the mite away and therefore not laying eggs on them. These reports agree with the work done by Wesonga, *et al.* (2005) who reported that, the wild *Lycopersicon* accessions (*L. hirsutum* and *L. peruvianum*) were found to be the most repellent to the spider mites.

Table 1(a): Effect of tomato varieties on duration (days) of incubation, larvae and protonymph stages of Red spider mite (*T. evansi*)

Variety	No. of Eggs	Incubation period		Larvae		Protonymph	
		Duration (mean no of days)	Total no	Duration (Mean no of days)	Total no	Duration (Mean no of days)	
Cal J	20	4.1a	18	2.8a	13	1.7ab	
Onyx	20	4.1a	19	2.7a	14	1.7ab	
Roma VFN	20	4.2a	18	2.8a	13	2.0ab	
Riogrande	20	4.1a	20	2.8a	16	2.0ab	
Money maker	20	4.1a	20	2.8a	17	2.3a	
Eden F1	20	4.3a	20	2.8a	16	2.1ab	
Anna F1	20	4.2a	20	2.7a	14	1.7ab	
Wild type	20	4.3a	18	2.6a	10	1.0b	
Mean		4.18		2.76		1.83	
SE		0.30		0.14		0.15	
CV %		22.4		16.37		25.0	

Means in the same column followed by the same letter are not significantly different using Tukey HSD test, $\alpha = 0.05$.

Table 1(b): Effect of tomato varieties on duration (days) of detonymph stage, a dult stage, sex and mortality of Red spider mite (*T. evansi*).

Variety	Deutonymph		Adult		Sex Male	Sex Female	Mortality (%) (Egg to Adult)
	Total no	Duration (Mean no of days)	Total no	Total duration (Egg to Adult)			
Cal J	7	1.3ab	5	9.9a	1	4	75
Onyx	7	1.8a	4	10.3a	2	2	80
Roma VFN	5	1.3ab	3	10.3a	-	3	85
Riogrande	12	1.5ab	9	10.4a	2	7	55
Money maker	13	1.8a	10	11.0a	3	7	50
Eden F1	9	1.3ab	5	10.5a	2	3	75
Anna F1	9	1.3ab	6	9.9a	2	4	70
Wild type		1.0b	1	8.9b	-	1	95
Mean		1.41		10.17			
SE		0.14					
CV %		31.33					

Means in the same column followed by the same letter are not significantly different using Tukey HSD test, $\alpha = 0.05$.

Table 2: Effect of tomato varieties on fecundity and longevity of *T. evansi*

Variety	Fecundity (Mean no. of eggs laid /female)	Average Longevity days
Cal J	32.0b	7.5b
Onyx	23.0c	7.2b
Roma VFN	23.9c	6.7b
Riogrande	48.6a	8.2b
Money maker	52.4a	12.8a
Eden F1	31.3b	7.6b
Anna	33.2b	7.4b
Wild type	9.1d	4.8c
Mean	31.94	7.78
SE	1.48	0.34
CV %	19.03	13.8

Means in the same column followed by the same letter are not significantly different using Tukey HSD test, $\alpha = 0.05$

V. CONCLUSION

The development, fecundity, adult longevity and survival of *T. evansi* varied with varieties. It was evident that antibiotic effect existed especially in Wild tomato variety. Tomato variety Anna F1 had higher yields in spite of high mite population and damages an indication of tolerance to mite damage. The present study did not study the mechanisms of resistance and there is a need to comprehensively study the mechanisms and sources of resistance.


VI. ACKNOWLEDGEMENT

This work was funded by Egerton University, Research and extension department. The authors wish to thank the following: Director Kenya Agricultural Research Institute (KARI), Njoro, Kenya, for material and facility support. Mr. Kamundia of KARI, Njoro Kenya, for statistical analysis and specialist advice on experimental design. The department of Crop, Horticulture and Soil sciences of Egerton University for their critical comments and guidance during research proposal development.

REFERENCES

- Cedola VC, Sanchez NE (2003) Effect of tomato pubescence on development, survival and fecundity of *Tetranychus urticae* Koch and *Neoseiulus californicus* (McGregor) (Acari: Tetranychidae, Phytoseiidae). *Acarologia* 3:255-260.
- Chatzivasilieadis E A, Boon J J, Sabelis M W (1999) Accumulation and turnover of 2-tridecanone in *Tetranychus urticae* and its consequences for resistance of wild and cultivated tomatoes. *Experimental and Applied Acarology* 23:1011-1021.
- Chatzivasilieadis E A, Sabelis M W (1997) Toxicity of methyl ketons from tomato trichomes to *Tetranychus urticae* Koch. *Experimental and Applied Acarology* 21:473-484.
- Chatzivasilieadis E A, Sabelis M W (1998) Variability in susceptibility among cucumber and tomato strains of *Tetranychus urticae* Koch to 2-tridecanone from tomato trichomes: Effects of host plant shift. *Experimental and Applied Acarology* 22:455-466.
- FAO (2001) State of food and agriculture. Food and Agriculture Organization. Rome, Italy.
- Helle W, Sabelis M W (1985) Spider mites, their biology, natural enemies and control, Vol. 1 B Elsevier, Amsterdam: 141-160.
- Isutsa D K, Wesonga J M, Onyango M O, Njoroge J B, Waiganjo M M, Agong S G, Fricke A, Hau B, Stuetzel H (2006) Proceedings of the Fifth Workshop on Sustainable Horticultural Production in the Tropics, Egerton University, Njoro, Kenya.
- Kamau A W (1985) The biology and control of tomato russet mite, *Acolops Lycopersci* (Masse) (Acarina: Eryophyidae) in Kenya, PhD Thesis, University of Nairobi.
- KARI (1996). Kenya Agricultural Research Institute Annual report Kennedy G G 2003. Tomato, pests parasitoids and predators: titrophic interactions involving the genus *Lycopersicon*. *Annual Review of Entomology* 48:51-72
- Maluf W R, Campos G A, Das Gracias C M (2001) Relationships between trichome types and spider mites (*Tetranychus evansi*) repellence in tomatoes with respect foliar Zingiberence Contents. *Euphytica*. 121 (1). 73-80.
- Meyer M K (1996) Mite pests and their predators on cultivated plants in South Africa. Vegetables and berries. Plant protection research institute handbook No. 6, ARC – Plant Protection research Institute. Pretoria.
- Murungi L K, Knapp M, Masinde P W, Onyambu G, Gitonga L, Agong S G (2009) Host-Plant acceptance, fecundity and longevity of *Tetranychus evansi* (Acari:Tetranychidae) on selected tomato accessions. Jomo Kenyatta university of Agriculture and Technology. Nairobi Kenya.
- Nihoul P (1993) Controlling glass house climate influences the interaction between tomato glandular trichome spider mite and predatory mite. *Crop protection* 12:443-447
- Quereshi A H, Oatman E R, Fleschner C A (1969) Biology of the spider mite, *Tetranychus evansi*. *Annals of Entomological society of America* 62:899-902.
- Ramalho F S, Fletchmann C H (1997) Niveis de infestacao de *Tetranychus evansi* Baker and Pritchard, 1960 em diferentes fases de desenvolvimento do tomateiro. *Rev. Agricult.* 54, 51-56.
- SAS Institute (2001) SAS/SAT, Version 8.02.SAS Institute Inc; Cary, NC., USA.
- Saunyama I G, Knapp M (2003).The effect of pruning and trellising of tomatoes (*Lycopersicon esculentum* Mill) on red spider mite (*Tetranychus evansi* Baker and Pritchard) incidence and crop yield in Zimbabwe. *Afri.crop Sci.J.* 11, 269-277.
- Simmons A T, Gurr G M (2005) Trichomes of *Lycopersicon* species and their hybrids:effect on pests and natural enemies. *Agricultural and Forest entomology* 7:265-276.
- Snyder J C, Carter C D (1984) Leaf trichomes and resistance of *Lycopersicon hirsutum* and *L. esculentum* to spider mites *Tetranychus urticae* Koch. *J.Amer.soc.Hort.Sci.* 109:837-843.
- Snyder J C, Carter C D, (1985) Trichomes on leaves of *Lycopersicon hirsutum* and *L. esculentum* and their hybrids. *Euphytica* 34: 53-64.
- Van Haren R, Steenhuis M, Sabelis M W, De ponti O (1987) Tomato stem trichomes and dispersal success of *Phytoseiulus persimilis* relative to its prey *Tetranychus urticae* .*Experimental and applied Acarology* 3:115-121.
- Varela A A, Seif A A, Loehr B (2003) A guide to IPM in tomato production in Eastern and Southern Africa. ICIPE Science Press, Nairobi. PP 21-26.
- Wesonga J M, Masinde P W, Losenge T Ngamau K, Agong S G, Anjichi V, Fricke A, Hau B, Stuetzel H (2005) Proceedings of the Fourth Workshop on Sustainable Horticultural Production in the Tropics. Department of Horticulture, Moi University, Eldoret, Kenya.
- Yang P M, Les S, Y, Joliffe F (2004) Farmers knowledge perceptions and practices in transgenic Bt Cotton in small producer systems in northern China. *Crop Protection Journal*.

PERSONAL PROFILE

	Name	: Mr. Musah, Samuel Matika
	Address	: P. O. Box 14197-20100, Nakuru, Kenya
	Mobile	: 0717 691 333 / 0723 884 139
	E-mail	: msmatika@yahoo.com
	Date of birth	: 05/09/1975 (Kakamega County, Kenya)
	Nationality	: Kenyan

EDUCATION

Year	Degree	Institution
2013	PhD Plant Pathology (In progress)	Karatina University, Kenya
2012	MSc Agronomy	Egerton University, Kenya
2002	BSc Agriculture Education and Extension	Egerton University, Kenya
1994	Kenya Certificate of Secondary Education	Musingu School, Kakamega
1990	Kenya Certificate of Primary Education	Luanda Pri. School, Kakamega

PROFESSIONAL QUALIFICATIONS

Dates/Year	Title of Certificate	Institution Awarding
Sept 2011	Improving Liriomyza leaf miners and Thrips monitoring and management in vegetables cropping systems for the Eastern African countries.	International centre for insect Physiology and Ecology (ICIPE, Kenya)
June 2012	Socio-Economic Approach to Research, Extension and Training.	Ministry of Agriculture, Kenya
March 2012	Participatory Analysis of Poverty and Livelihood Dynamics (PAPOLD)	Ministry of Agriculture, Kenya
June 2011	Positive Management	African Virtual University

FIELD OF SPECIALIZATION

Agronomy, Plant Pathology, Agriculture Education and Extension
--

WORK EXPERIENCE

Apr2012- Present	Part time Lecturer, Karatina University, Laikipia University and University of Kabianga.
Jun'08 - Present	Chief Agriculture Extension Officer, Nakuru County, Kenya
Jun'06 – Jun'08	Graduate Teacher, Teachers Service Commission of Kenya
Mar 2003- Apr2004	Production Manager, Migotiyu Plantations ltd, Kenya

RECENT RESEARCH PROJECT

Project period	Research Topic
Jan2009 –Dec2011	Distribution and the use of varietal resistance and pesticide for the control of Red spider mite <i>Tetranychus evansi</i> (Baker & Pritchard) in tomato <i>Lycopersicon esculentum</i> (Mill), Kenya. Donor: Egerton University, Kenya. Research and Extension Department (Principle Investigator: Musah Samuel Matika)

RECENT PUBLICATION AND SCIENTIFIC CONFERENCE ATTENDED

1	Musah S.M., Kamau A.W., Munene M. Use of varietal resistance in the control of red spider mite <i>Tetranychus evansi</i> (Baker & pritchard) in tomato <i>Lycopersicon esculentum</i> (mill) production in Kenya. Conference paper presented in Jaramogi Oginga Odinga University of Science and Technology, First International Research Conference. Dates: 24th – 26 th June 2015 Main Campus, Bondo, Kenya
2	Musah S.M., Kamau A.W., Munene M. Effect of Pesticides on the Control of Red Spider Mite <i>Tetranychus evansi</i> (Backer & Pritchard) on Tomato <i>Lycopersicon esculentum</i> (Mill).Conference paper presented in Chuka University, Second International Research Conference. Dates: 28th – 30 th October 2015 Main Campus, Chuka University, Kenya.
3	Musah S.M., Kamau A.W., Munene M. Survey on Distribution and Damage on Tomatoes by Red Spider Mite in Subukia and Rongai Sub County, Kenya.Conference paper presented in Chuka University, Second International Research Conference. Dates: 28th – 30 th October 2015 Main Campus, Chuka University, Kenya.
4	Mulehi, H., Wawire, A., Musah, S.M. Evaluation of E-Extension in Agriculture: The Case of Molo Sub-County, in Nakuru County, Kenya. Conference paper presented in Chuka University, Second International Research Conference. Dates: 28th – 30 th October 2015 Main Campus, Chuka University, Kenya.

MEMBERSHIP TO PROFESSIONAL / SCIENTIFIC ASSOCIATIONS

Membership Category	Organization
Member	Agricultural Professionals Society of Kenya
Member	Entomological Society of East Africa
Member	Teachers Service Commission (TSC NO 473947)
Member	Society for Economic Botany-Kenya Chapter